

Exact Charge Dynamics in the Supersymmetric t - J Model with Inverse-Square Interaction

Mitsuhiro Arikawa,¹ Takashi Yamamoto,² Yasuhiro Saiga,³ and Yoshio Kuramoto¹

¹*Department of Physics, Tohoku University, Sendai 980-8578, Japan*

²*Max-Planck-Institut für Physik komplexer Systeme, Nöthnitzer Str. 38, D-01187 Dresden, Germany*

³*Institute for Solid State Physics, University of Tokyo, Roppongi 7-22-1, Tokyo 106-8666, Japan*

Dynamical charge structure factor $N(Q, \omega)$ with Q smaller than the Fermi wave number is derived analytically for the one-dimensional supersymmetric t - J model with $1/r^2$ interaction. Strong spin-charge separation in dynamics is found. $N(Q, \omega)$ with given electron density does not depend on the spin polarization for arbitrary size of the system. In the thermodynamic limit, only two holons and one antiholon contribute to $N(Q, \omega)$. These results together with generalized ones for the $SU(K,1)$ supersymmetric t - J model are derived via solution of the Sutherland model in the strong coupling limit.

75.10.Jm, 05.30.-d, 71.27.+a

Strong correlation between one-dimensional electrons has drastic consequences to their physical properties. The most spectacular is the spin-charge separation; the elementary excitations are spinons and holons both of which obey the fractional statistics [1]. This is in contrast with the fermionic Landau quasi-particles realized in higher dimensions. In the supersymmetric t - J model with $1/r^2$ interaction [2], these fractional particles appear in the simplest manner. Exact thermodynamics for the model [3,4] can be interpreted in terms of free spinons and holons. The charge susceptibility, for example, depends on the electron density, but is independent of the spin polarization. This kind of independence is referred to as the strong spin-charge separation [3,4]. It is natural to ask how the spin-charge separation appears in dynamical correlation functions.

In the absence of charge degrees of freedom, the supersymmetric t - J model is reduced to the spin chain called the Haldane-Shastry model [5,6]. For the latter model Haldane and Zirnbaier [7] derived exactly the dynamical structure factor which is non-zero only over a compact region in the momentum-frequency plane. Such region is called the support. For the supersymmetric t - J model, Ha and Haldane [8] analyzed numerical results for finite-size systems, and found that only a few number of elementary excitations contribute to spectral functions, namely dynamical structure factors. They proposed a compact support for each spectral function in the thermodynamic limit, but did not obtain the spectral functions themselves. Analytical knowledge should obviously provide deeper insight into strong correlation effects on dynamics. Recently, Kato partially derived a spectral function corresponding to the removal of a particle from the high-density limit [9]. So far all analytical work in the literature does not include charge degrees of freedom in the ground state, and derivation of charge dynamics has remained a theoretical challenge.

In this Letter, we report on exact analytical results on

charge dynamics for the one-dimensional supersymmetric t - J model with $1/r^2$ interaction at zero temperature, and its generalization from $SU(2,1)$ to $SU(K,1)$ supersymmetry. We show explicitly how the strong spin-charge separation manifests in charge dynamics. The charge correlation function is relevant to dielectric and elastic responses of real systems with quasi-one-dimensional electrons. Our work should motivate further analytical work toward other dynamical correlation functions which are relevant to various spectroscopies such as neutron scattering and photoemission.

We consider the supersymmetric t - J model given by

$$\mathcal{H}_{tJ} = \mathcal{P} \sum_{i < j} \left[-t_{ij} \sum_{\sigma=\uparrow, \downarrow} \left(c_{i\sigma}^\dagger c_{j\sigma} + h.c. \right) + J_{ij} \left(\mathbf{S}_i \cdot \mathbf{S}_j - \frac{1}{4} n_i n_j \right) \right] \mathcal{P}, \quad (1)$$

where $c_{i\sigma}$ is the annihilation operator of an electron at site i with spin σ , n_i is the number operator and $\mathbf{S}_i$ is the spin operator. The projection operator $\mathcal{P}$ excludes double occupation at each site. The transfer energy t_{ij} and exchange one J_{ij} are given by $t_{ij} = J_{ij}/2 = t D_{ij}^{-2}$ where $D_{ij} = (N/\pi) \sin(\pi(i-j)/N)$ with N being the number of lattice sites. This model is generalized to the $SU(K,1)$ supersymmetry as

$$\mathcal{H}_{tJ}^{(K)} = t \sum_{i < j} D_{ij}^{-2} \tilde{P}_{ij}. \quad (2)$$

Here we have introduced a graded permutation operator $\tilde{P}_{ij} = \sum_{\alpha, \beta} X_i^{\alpha\beta} X_j^{\beta\alpha} \theta_\beta$, where $X_j^{\beta\alpha}$ changes a state α to β at site j , with α, β being either 0 (hole state), or one of $\sigma = 1, \dots, K$ (spin state). The sign factor θ_β is -1 if $\beta = 0$ and 1 otherwise. In the case of $K = 2$ the expression (2) is reduced to eq.(1).

In terms of the density fluctuation operator $n_Q = N^{-1/2} \sum_{\sigma=1}^K \sum_{l=1}^N X_l^{\sigma\sigma} \exp(iQl)$ with $Q > 0$, the dynamical charge structure factor is defined by

$$N(Q, \omega) = \sum_{\nu} |\langle \nu | n_Q | 0 \rangle|^2 \delta(\omega - E_{\nu} + E_0), \quad (3)$$

where $|\nu\rangle$ denotes a normalized eigenstate of the Hamiltonian (2) with energy E_{ν} .

In the $SU(K,1)$ t - J model, a holon has the spectrum $\epsilon_h(q) = Ktq[2\pi(1-\bar{n})/K + q]/2$ with $\bar{n}$ being the average electron number per site, and another charge excitation called the antiholon has the spectrum $\epsilon_a(q) = tq[2\pi(1-\bar{n}) - q]/2$ [4,8]. The latter represents the excitation from the condensate of holons. We first give our main result of this Letter, and then its derivation. The system has N_{σ} electrons with spin σ . In the small momentum region $0 < Q \leq k_F^{\text{Min}}$, with k_F^{Min} being the smallest Fermi number of electrons, *i.e.*, $k_F^{\text{Min}} = \pi \text{Min}(N_1, \dots, N_K)/N$, we derive $N(Q, \omega)$ in the form:

$$N(Q, \omega) = A_0 Q^2 \int_0^{2\pi(1-\bar{n})} dq' \prod_{i=1}^K \int_0^{2\pi/K} dq_i \times \delta\left(Q - q' - \sum_{i=1}^K q_i\right) \delta\left(\omega - \epsilon_a(q') - \sum_{i=1}^K \epsilon_h(q_i)\right) \times \frac{\prod_{i < j} |q_i - q_j|^{2/K}}{\prod_{i=1}^K (Kq_i + q')^2} \epsilon_a(q')^{K-1} \prod_{i=1}^K \epsilon_h(q_i)^{1/K-1}, \quad (4)$$

where $A_0 = K/(2\pi) \prod_{j=1}^K \Gamma(1/K)/\Gamma(j/K)^2$. Surprisingly this $N(Q, \omega)$ has almost the same form as that of the spinless Sutherland model with the coupling parameter $1/K$ [10,11]. In the latter model excitations consist of K particles created above the pseudo Fermi level with charge $-e$, and a hole with charge Ke . The only difference in $N(Q, \omega)$ between the two models is the integration range of momenta. Thus the result above can be interpreted in terms of the fractional statistics [12].

With $K = 2$, it can be shown that $N(Q, \omega)$ diverges as $[\epsilon_h(Q) - \omega]^{-1/2}$ as the frequency approaches the upper edge corresponding to the holon dispersion [13]. The static structure factor $N(Q) = \int d\omega N(Q, \omega)$ reproduces the known result [14] obtained for the Gutzwiller wave function. This should be so since the exact ground state of the t - J model is given by the Gutzwiller wave function [2].

As in the previous study of thermodynamics [3,4,15,16], we use the Sutherland model with the $SU(K,1)$ internal symmetry as an auxiliary, and take the limit $\beta \rightarrow \infty$ of the coupling parameter β . Then the particles crystallize with equal distance from each other, and the resultant dynamics excluding phonons and uniform motion of the center of gravity is mapped to that of the $SU(K,1)$ t - J model given by eq.(2). The merit of using the Sutherland model is that much more is known about properties of the eigenfunctions than those for the t - J model. The Sutherland model with arbitrary internal symmetry is written in the form

$$\mathcal{H} = -\frac{1}{2M} \sum_{i=1}^N \frac{\partial^2}{\partial x_i^2} + \frac{\pi^2}{L^2 M} \sum_{i < j} \frac{\beta(\beta - s_{ij})}{\sin^2[\pi(x_i - x_j)/L]}, \quad (5)$$

where L is the size of the system, and s_{ij} represents the exchange operator of coordinates of particles i and j [17]. In order to reproduce the lattice model we take the limit of large β and M , keeping the ratio $t = \beta/M$ fixed. Here we take the lattice parameter L/N as the unit of length. Note that the symmetry of the wave function leads to the relation $s_{ij} = -\hat{P}_{ij}$ within the Hilbert space of identical particles.

Let us describe derivation of $N(Q, \omega)$ taking the $SU(2,1)$ Sutherland model. Generalization to the $SU(K,1)$ supersymmetry is straightforward. The system has N_h holes, $N_{\uparrow}$ up-spin electrons and $N_{\downarrow}$ down-spin ones, whose coordinates are represented by x_i^h for i -th hole, $x_i^{\uparrow}$ for i -th up-spin electron and $x_i^{\downarrow}$ for i -th down-spin electron. We arrange them as $x \equiv (x_1, x_2, \dots, x_N) = (x_1^h, \dots, x_{N_h}^h, x_1^{\uparrow}, \dots, x_{N_{\uparrow}}^{\uparrow}, x_1^{\downarrow}, \dots, x_{N_{\downarrow}}^{\downarrow}) \equiv (x^h, x^{\uparrow}, x^{\downarrow})$ and assume that N_h is even, and both $N_{\uparrow}$ and $N_{\downarrow}$ are odd. Then the ground state is non-degenerate. The interval $I = [1, N]$ denotes $\{i \in \mathbf{Z} | 1 \leq i \leq N\}$ for a positive integer N . We define $I_h = [1, N_h]$, $I_{\uparrow} = [N_h + 1, N_h + N_{\uparrow}]$ and $I_{\downarrow} = [N_h + N_{\uparrow} + 1, N]$. Without loss of generality we assume $N_{\downarrow} \geq N_{\uparrow}$. The wave function of the ground state for a set of $N_h, N_{\uparrow}, N_{\downarrow}$ is given by,

$$\Psi_{\text{GS}} = \prod_{i < j \in I} \left(1 - \frac{z_j}{z_i}\right)^{\beta} \prod_{\sigma=\uparrow, \downarrow} \prod_{i < j \in I_{\sigma}} \left(1 - \frac{z_j}{z_i}\right), \quad (6)$$

where the complex coordinates $z = (z_1, \dots, z_N)$ are related to the original ones $x = (x_1, \dots, x_N)$ by $z_j = \exp(2\pi i x_j / L)$.

The spectrum of the Sutherland model is conveniently analyzed with use of a similarity transformation generated by $\mathcal{O} = \prod_{i < j \in I} (1 - z_j/z_i)^{\beta} \prod_{i \in I} z_i^{(N_{\downarrow}-1)/2}$. The transformed Hamiltonian $\hat{\mathcal{H}} = \mathcal{O}^{-1} \mathcal{H} \mathcal{O} - e_0$ describes the excited states relative to the spinless particle model with the ground-state energy $e_0 = \pi^2 \beta^2 N(N-1)/6ML^2$. We obtain

$$\hat{\mathcal{H}} = \frac{1}{2M} \left(\frac{2\pi}{L}\right)^2 \sum_{i=1}^N \left(\hat{d}_i + \beta \frac{N-1}{2} - \frac{N_{\downarrow}-1}{2}\right)^2. \quad (7)$$

Here $\hat{d}_i$ is called the Cherednik-Dunkl operator [18] and is given by

$$\hat{d}_i = z_i \frac{\partial}{\partial z_i} + \beta \sum_{k < i} \frac{z_i}{z_i - z_k} (1 - s_{ik}) + \beta \sum_{i < k} \frac{z_k}{z_i - z_k} (1 - s_{ik}) + \beta(1 - i). \quad (8)$$

It is known that $\hat{d}_i$ can be diagonalized simultaneously with real eigenvalue $\bar{\lambda}_i$. The resultant eigenfunction are

polynomials $E_\lambda(z; \beta)$ of z_i such as $\prod_i z_i^{\lambda_i} + \dots$, and are called non-symmetric Jack polynomials [18]. Since we are dealing with identical particles, eigenfunction should satisfy the following conditions of the $SU(2,1)$ supersymmetry:

- (i) symmetric with respect to exchange between z_i^h 's;
- (ii) anti-symmetric with respect to exchange between z_i^σ 's with the same σ .

By taking linear combination of $E_\lambda(z; \beta)$, we can construct a polynomial $K_\lambda(z; \beta)$ with the $SU(2,1)$ supersymmetry. The latter belongs to a family called the Jack polynomial with prescribed symmetry [19–21]. We specify the set of momenta as $(\lambda^h, \lambda^\uparrow, \lambda^\downarrow) \equiv (\lambda_1^h, \dots, \lambda_{N_h}^h, \lambda_1^\uparrow, \dots, \lambda_{N_\uparrow}^\uparrow, \lambda_1^\downarrow, \dots, \lambda_{N_\downarrow}^\downarrow)$ with $\lambda_1^h \geq \dots \geq \lambda_{N_h}^h, \lambda_1^\uparrow > \dots > \lambda_{N_\uparrow}^\uparrow$ ($\sigma = \uparrow, \downarrow$). In this way we can parameterize $K_\lambda(z; \beta)$ by using $\lambda \equiv (\lambda^h, \lambda^\uparrow, \lambda^\downarrow)$. At the ground state we have $\lambda = \lambda_{GS} = (\mu^h, \mu^\uparrow, \mu^\downarrow)$ with $\mu^h = ((N_\downarrow - 1)/2, (N_\downarrow - 1)/2, \dots, (N_\downarrow - 1)/2)$, $\mu^\uparrow = ((N_\uparrow + N_\downarrow)/2 - 1, (N_\uparrow + N_\downarrow)/2 - 2, \dots, (N_\downarrow - N_\uparrow)/2)$ and $\mu^\downarrow = (N_\downarrow - 1, N_\downarrow - 2, \dots, 0)$.

We define the inner product for functions $f(z)$ and $g(z)$ in N -complex variable $z = (z_1, \dots, z_N)$ as follows:

$$\langle f, g \rangle_0 = \prod_{j=1}^N \oint_{|z_j|=1} \frac{dz_j}{2\pi i z_j} \overline{f(z)} g(z) \prod_{k < l \in I} |z_k - z_l|^{2\beta}, \quad (9)$$

where $\overline{f(z)}$ denotes the complex conjugation of $f(z)$. As is clear from the definition of the transformation $\mathcal{O}$, $K_\lambda(z; \beta)$ is orthogonal with respect to the above inner product. In order to derive the norm of $K_\lambda(z; \beta)$, we generalize the procedure taken in refs. [21–23]. The result of lengthy calculation is given by

$$\langle K_\lambda, K_\lambda \rangle_0 = N_h! N_\downarrow! N_\uparrow! \rho_\lambda^{-1} \langle E_\lambda, E_\lambda \rangle_0, \quad (10)$$

where $\langle E_\lambda, E_\lambda \rangle_0$ denotes the norm of the non-symmetric Jack polynomials. We refer to the literature [18] for the explicit form of the norm since it requires many lengthy combinatorial quantities. The quantity $\rho_\lambda = \rho_\lambda^h \rho_\lambda^\uparrow \rho_\lambda^\downarrow$ is given by

$$\rho_\lambda^h = \prod_{i < j \in I_h} \frac{\bar{\lambda}_i - \bar{\lambda}_j + \beta}{\bar{\lambda}_i - \bar{\lambda}_j}, \quad \rho_\lambda^\sigma = \prod_{i < j \in I_\sigma} \frac{\bar{\lambda}_i - \bar{\lambda}_j - \beta}{\bar{\lambda}_i - \bar{\lambda}_j}. \quad (11)$$

In the lattice model, the density correlation of particles is the same as that of holes because of the completeness relation $\sum_{\sigma=\uparrow, \downarrow} X_i^{\sigma\sigma} + X_i^{00} = 1$. In the Sutherland model, on the other hand, the correlation function of the sum of hole-density and particle-density gives the phonon spectrum. It can be shown that the intensity of the phonon correlation function is smaller than the spin correlation function by a factor of $O(\beta^{-1})$. In the leading order in β^{-1} , the correlation function of holes is the same as that of particles. In view of this situation we consider the power sum $p_m = \sum_{i \in I_h} z_i^m$ for holes. The

hole-density n_Q can be expressed as $n_Q = p_m / \sqrt{N}$ with $Q = 2\pi m / N$ [10, 11]. In order to calculate $N(Q, \omega)$, we need to know the expansion coefficient c_λ which appears in

$$p_m \Psi_{GS} = \sum_\lambda c_\lambda K_\lambda(z; \beta) \prod_{i < j \in I} \left(1 - \frac{z_j}{z_i}\right)^\beta. \quad (12)$$

It is difficult to derive c_λ for general values of m . However, there occurs drastic simplification for the small momentum region with $m \leq (N_\uparrow - 1)/2$. In this region one can show that $p_m \Psi_{GS}$ has only charge excitations due to the triangular structure of the polynomial $K_\lambda(z; \beta)$ [24].

Without spin excitations, we have the explicit expression of $K_\lambda(z; \beta)$ following Baker and Forrester [19] as follows:

$$K_\lambda(z; \beta) = J_\nu^\alpha(z_h) \prod_{i \in I} z_i^{(N_\downarrow - 1)/2} \prod_{\sigma=\uparrow, \downarrow} \prod_{k < l \in I_\sigma} \left(1 - \frac{z_l}{z_k}\right), \quad (13)$$

where $\alpha = 2 + 1/\beta$ and $J_\nu^\alpha(z_h) = J_\nu^\alpha(z_1, \dots, z_{N_h})$ is the symmetric Jack polynomial [18]. The charge excitations are characterized by the set $\nu = (\nu_1, \dots, \nu_{N_h})$, which is related to λ by $\lambda = (\nu, 0, 0) + \lambda_{GS}$.

Since p_m can be expanded by the symmetric Jack polynomials [25], we can determine the coefficient c_λ in eq.(12). With use of a square $s = (i, j)$ in the Young diagram ν , we introduce the notations

$$[0']_\nu^\alpha = \prod_{\substack{s \in \nu \\ s \neq (1,1)}} [\alpha a'_\nu(s) - l'_\nu(s)], \quad h_\nu^\alpha = \prod_{s \in \nu} [\alpha(a_\nu(s) + 1) + l_\nu(s)],$$

where $a_\nu(s) = \nu_i - j$ is the arm-length, $a'_\nu(s) = j - 1$ is the arm-colength, $l_\nu(s) = \nu'_j - i$ with ν'_j the length of the column j is the leg-length, and $l'_\nu(s) = i - 1$ is the leg-colength. We then obtain the dynamical structure factor of the Sutherland model with the coupling parameter β as follows:

$$N(Q, \omega; \beta) = \frac{\alpha^2 m^2}{N} \sum_\nu \delta_{m, |\nu|} \delta(\omega - E_\lambda + E_{\lambda_{GS}}) \times \frac{\langle K_\lambda, K_\lambda \rangle_0}{\langle K_{\lambda_{GS}}, K_{\lambda_{GS}} \rangle_0} \left(\frac{[0']_\nu^\alpha}{h_\nu^\alpha} \right)^2, \quad (14)$$

where $Q = 2\pi m / N$ and $|\nu| = \sum_{j=1}^{N_h} \nu_j$. The excitation energy is given by

$$E_\lambda - E_{\lambda_{GS}} = \frac{\beta}{2M} \sum_{i=1}^{N_h} [\alpha \nu_i^2 + (N_h + 1 - 2i) \nu_i]. \quad (15)$$

By taking the limit $\beta \rightarrow \infty$ with the ratio $t = \beta / M$ kept fixed, we can derive $N(Q, \omega)$ of the t - J model (1). In this limit $N(Q, \omega)$ simplifies dramatically by the following fact: The coefficient $[0']_\nu^\alpha$ in eq.(14) vanishes if the partition ν contains $s = (3, 2)$. Then $N(Q, \omega)$ is

determined by three parameters which are related directly with momentum of the elementary excitations: two holons and one antiholon. We emphasize that $N(Q, \omega)$ does not depend on the spin polarization as long as Q is smaller than the smallest Fermi wave number of all electrons.

Derivation of dynamical charge structure factor for $SU(K, 1)$ model proceeds in a similar manner. We write the momenta of K holons and one antiholon by $\lambda_1, \dots, \lambda_K$ and λ' , respectively. Then $N(Q, \omega)$ for finite N and arbitrary K can be expressed as follows:

$$\begin{aligned}
N(Q, \omega) = & B_0 m^2 \sum_{\lambda_1 \geq \dots \geq \lambda_K, \lambda'} \delta_{m, ||\lambda||} \delta(\omega - E_{\lambda\lambda'}) \\
& \times \frac{\Gamma(\lambda') \Gamma(N_h - \lambda' + K)}{\Gamma(\lambda' + 1 - K) \Gamma(N_h - \lambda' + 1)} \\
& \times \prod_{j=1}^K \frac{1}{(K\lambda_j + \lambda' - j)(K(\lambda_j - 1) + \lambda' - (j - 1))} \\
& \times \prod_{j=1}^K \frac{\Gamma(\lambda_j - (j - 1)/K) \Gamma(\lambda_j + (N_h - j + 1)/K)}{\Gamma(\lambda_j - (j - K)/K) \Gamma(\lambda_j + (N_h - j + K)/K)} \\
& \times \prod_{1 \leq i < j \leq K} \left[\frac{\Gamma(\lambda_i - \lambda_j + 1 - (i - j)/K)}{\Gamma(\lambda_i - \lambda_j + 1 - (i - j + 1)/K)} \right. \\
& \times \left. \frac{\Gamma(\lambda_i - \lambda_j - (i - j - 1)/K)}{\Gamma(\lambda_i - \lambda_j - (i - j)/K)} \right], \quad (16)
\end{aligned}$$

where $||\lambda|| = \lambda' + \sum_{j=1}^K (\lambda_j - 1)$ and $B_0 = 2\pi A_0 K^{1-K} K! / N$. The excitation energy is given by

$$\begin{aligned}
E_{\lambda\lambda'} = & \frac{t}{2} \left(\frac{2\pi}{N} \right)^2 \{ \lambda' (N_h + K - \lambda') \\
& + \sum_{j=1}^K [(\lambda_j - 1)(N_h + K(\lambda_j + 1) - (2j - 1))] \}. \quad (17)
\end{aligned}$$

In the special case of $\lambda_K = 0$ where less than K holons are excited, we need to modify the form of eq.(16). However, this case can be neglected in the thermodynamic limit. We have checked the validity of eq.(16) together with the modified one in the case of $K = 2$ by comparison with numerical results up to $N = 16$.

Following the same procedure as that in the spinless Sutherland model [10,11], we obtain the expression (4) in the thermodynamic limit. The present derivation makes it clear that the spin-charge separation persists in systems with finite size. In the $SU(K, 1)$ model, α in eq.(13) is given by $\alpha = K + \beta^{-1}$, which tends to K in the limit of $\beta \rightarrow \infty$. Since $1/\alpha$ can be regarded as the coupling constant in the charge sector of the fractional statistics,

$N(Q, \omega)$ given by eq.(4) has the apparent coupling constant $1/K$.

The authors would like to thank Y. Kato for valuable discussions. T.Y. and Y.S. wish to acknowledge the support of the CREST from the Japan Science and Technology Corporation. T.Y. wishes to thank the support of the Visitor Program of the MPI-PKS.

-
- [1] F.D.M. Haldane, Phys. Rev. Lett. **66**, 1529 (1991).
 - [2] Y. Kuramoto and H. Yokoyama, Phys. Rev. Lett. **67**, 1338 (1991).
 - [3] Y. Kuramoto and Y. Kato, J. Phys. Soc. Jpn. **64**, 4518 (1995).
 - [4] Y. Kato and Y. Kuramoto, J. Phys. Soc. Jpn. **65**, 1622 (1996).
 - [5] F.D.M. Haldane, Phys. Rev. Lett. **60**, 635 (1988).
 - [6] B.S. Shastry, Phys. Rev. Lett. **60**, 639 (1988).
 - [7] F.D.M. Haldane and M.R. Zirnbauer, Phys. Rev. Lett. **71**, 4055 (1993).
 - [8] Z.N.C. Ha and F.D.M. Haldane, Phys. Rev. Lett. **73**, 2887 (1994).
 - [9] Y. Kato, Phys. Rev. Lett. **81**, 5402 (1998).
 - [10] Z.N.C. Ha, Phys. Rev. Lett. **73**, 1574 (1995); **74**, 620 (1995) (E); Nucl. Phys. **B 435**, 604 (1995).
 - [11] F. Lesage, V. Pasquier and D. Serban, Nucl. Phys. **B 435**, 585 (1995).
 - [12] Y. Kato, T. Yamamoto and M. Arikawa, J. Phys. Soc. Jpn. **66**, 1954 (1997).
 - [13] E.R. Mucciolo, B.S. Shastry, B.D. Simons and B.L. Altshuler, Phys. Rev. B **49**, 15197 (1994).
 - [14] F. Gebhard and D. Vollhardt, Phys. Rev. B **38**, 6911 (1988).
 - [15] A. Polychronakos, Phys. Rev. Lett. **70**, 2329 (1990).
 - [16] B. Sutherland and B.S. Shastry, Phys. Rev. Lett. **71**, 5 (1993).
 - [17] A. Polychronakos, Phys. Rev. Lett. **69**, 703 (1992).
 - [18] I.G. Macdonald, *Symmetric Functions and Orthogonal Polynomials*, (American Mathematical Society, 1997).
 - [19] T.H. Baker and P.J. Forrester, Nucl. Phys. **B 492**, 682 (1997).
 - [20] C.F. Dunkl, Comm. Math. Phys **197**, 451 (1998).
 - [21] Y. Kato and T. Yamamoto, J. Math. A: Math. Gen. **31**, 9171 (1998).
 - [22] K. Takemura and D. Uglov, J. Phys. A: Math. Gen. **30**, 3685 (1997).
 - [23] Y. Kato, private communication (1997).
 - [24] Y. Kato and Y. Kuramoto, Phys. Rev. Lett. **74**, 1222 (1995).
 - [25] P.J. Hanlon, R.P. Stanley and J.R. Stembridge, Contemp. Math. **138**, 151 (1992).